

Tablets

Composition

Ancozine Tablets

Each tablet contains:

Meclozine HCl 25 mg Pyridoxine HCl 50 mg

Action

Meclozine HCl is an H_1 histamine receptor blocker of the piperazine side chain group; it acts on CNS by its ability to block muscarinic receptors in the brain.

Pyridoxine HCl is a water-soluble vitamin involved in amino acid, carbohydrate, and fat metabolism. It is also required for the formation of haemoglobin. It is used to prevent uncontrolled excitation of CNS, and eventually uncontrolled muscle seizures.

This combination provides a synergistic anti-emetic effect.

Indications

Ancozine is an anti-emetic, effective in the management of nausea, vomiting, and dizziness during pregnancy or dizziness associated with motion sickness.

Ancozine has been found useful in the management of vertigo associated with diseases affecting the vestibular system.

Contraindications

Ancozine is contraindicated in patients hypersensitive to any of the components, in patients with asthma, glaucoma, chronic pulmonary disease, or any difficulty in urination.

Adverse Reactions

Ancozine may cause drowsiness, dry mouth, and blurred vision on rare occasions.

Warnings

Ancozine may cause drowsiness, so it is better not to operate dangerous machines during therapy.

Ancozine has a synergistic drowsiness effect with alcohol, sedatives, and tranquilizers.

Dosage and Administration

Pregnancy

The usual dose for nausea and vomiting during pregnancy is two tablets at night followed by one tablet in the morning if necessary.

Motion Sickness

The initial dose of one or two tablets should be taken one hour prior to embarkation for protection against motion sickness. Thereafter, the dose may be repeated every 24 hours for the rest of the journey.

Vertigo

For the control of vertigo associated with diseases affecting the vestibular system, the recommended dose is one to four tablets in divided doses, 2-3 times daily depending upon clinical response.

Presentation

Ancozine Tablets

Box of 20 tablets.